

Cayley transforms of $su(2)$ representations

T. S. Van Kortryk

120 Payne Street, Paris MO65275
vankortryk@gmail.com

Abstract

Cayley rational forms for rotations are given as explicit matrix polynomials for any quantized spin j . The results are compared to the Curtright-Fairlie-Zachos matrix polynomials for rotations represented as exponentials.

Curtright, Fairlie, and Zachos (CFZ) obtained an elegant and intuitive result [1] expressing the exponential form for a rotation about an axis $\hat{\mathbf{n}}$, for *any* quantized angular momentum j , as a polynomial of order $2j$ in the corresponding $(2j+1) \times (2j+1)$ spin matrices $\hat{\mathbf{n}} \cdot \mathbf{J}$, thereby culminating several decades of previous studies on this or closely related problems [2]-[6]. For each angle-dependent coefficient of the polynomial, the explicit formula found by CFZ involves nothing more complicated than a truncated series expansion for a power of the arcsin function. Although a detailed proof of the CFZ result is not exhibited in [1], the essential ingredients needed to provide such a proof are in that paper, and indeed, details of two elementary derivations were subsequently given in [7].

I discuss here the corresponding polynomial result for the Cayley rational form of an irreducible, unitary $SU(2)$ representation with spin j . In this case, the explicit polynomial coefficients involve nothing more complicated than truncated series expansions for simple, finite products. I compare the Cayley form to the CFZ result.

The CFZ formula for a rotation through an angle θ about an axis $\hat{\mathbf{n}}$, valid for any spin $j \in \{0, \frac{1}{2}, 1, \frac{3}{2}, \dots\}$, is given by the unitary $SU(2)$ matrix

$$U = \exp(i \theta \hat{\mathbf{n}} \cdot \mathbf{J}) = \sum_{k=0}^{2j} \frac{1}{k!} A_k^{[j]}(\theta) (2i \hat{\mathbf{n}} \cdot \mathbf{J})^k, \quad (1)$$

where the angle-dependent coefficients of the various spin matrix powers are explicitly given by

$$A_k^{[j]}(\theta) = \sin^k(\theta/2) (\cos(\theta/2))^{\epsilon(j,k)} \left[\frac{1}{(\sqrt{1-x})^{\epsilon(j,k)}} \left(\frac{\arcsin \sqrt{x}}{\sqrt{x}} \right)^k \right]_{x=\sin^2(\theta/2)}^{\text{Trunc}_{[j-k/2]}}. \quad (2)$$

Here, $\lfloor \dots \rfloor$ is the integer-valued floor function while $\epsilon(j, k)$ is a binary-valued function of $2j - k$ that distinguishes even and odd integers: $\epsilon(j, k) = 0$ for even $2j - k$, and $\epsilon(j, k) = 1$ for odd $2j - k$. More importantly, $\text{Trunc}_n[f(x)]$ is the n th-order Taylor polynomial truncation for any $f(x)$ admitting a power series representation:

$$f(x) = \sum_{m=0}^{\infty} f_m x^m, \quad \text{Trunc}_n[f(x)] \equiv \sum_{m=0}^n f_m x^m. \quad (3)$$

To emulate the CFZ formula, I now construct Cayley transforms [8, 9] of elements in the $su(2)$ algebra as spin matrix polynomials.

At first glance the Cayley transform for any spin representation would seem to follow immediately from the CFZ result just by changing variables from θ to a function of the angle, $\alpha(\theta)$. For any given numerical value of $\hat{\mathbf{n}} \cdot \mathbf{J}$ this would be so, of course, but it is not obviously so for matrix-valued $\hat{\mathbf{n}} \cdot \mathbf{J}$. In fact, it turns out that the matrix produced by the Cayley transform of an $su(2)$ generator reduces to a spin matrix polynomial that has a simpler form than the CFZ result (1) for the exponential of that generator.

For a finite dimensional, unitary, irreducible representation of $SU(2)$, all the underlying spin matrices $iS \equiv 2i \hat{\mathbf{n}} \cdot \mathbf{J}$ are anti-hermitian, and for spin j satisfy the Cayley-Hamilton theorem [8, 10] appropriate for $(2j+1) \times (2j+1)$ matrices. Consequently the Cayley rational form of a unitary $SU(2)$ group element for spin j can be reduced to a spin matrix polynomial [11]:

$$U = \frac{1 + 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J}}{1 - 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J}} = \sum_{k=0}^{2j} \mathfrak{A}_k^{[j]}(\alpha) (2i \hat{\mathbf{n}} \cdot \mathbf{J})^k, \quad (4)$$

where α is a real parameter, and where the coefficients $\mathfrak{A}_k^{[j]}(\alpha)$ are to be determined as functions of α . The challenge here is to rewrite the geometric series $1/(1 - 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J})$ for spin j as a polynomial in $\hat{\mathbf{n}} \cdot \mathbf{J}$. Thus define

$$\frac{1}{1 - 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J}} = \sum_{k=0}^{2j} \mathfrak{B}_k^{[j]}(\alpha) (2i \hat{\mathbf{n}} \cdot \mathbf{J})^k. \quad (5)$$

Then clearly $\mathfrak{A}_0^{[j]} = 2\mathfrak{B}_0^{[j]} - 1$ and $\mathfrak{A}_{k \geq 1}^{[j]} = 2\mathfrak{B}_{k \geq 1}^{[j]}$. The coefficients in the latter expansion follow directly from the methods in [1, 7]. The results are succinctly given by

$$\mathfrak{B}_k^{[j]}(\alpha) = \frac{\alpha^k}{\det(1 - 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J})} \text{Trunc}_{2j-k}[\det(1 - 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J})], \quad (6)$$

where the truncation is in powers of α , and where the determinant for spin j is

$$\det(1 - 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J}) = \prod_{m=0}^{2j} (1 - 2i\alpha(j-m)) = \prod_{n=1}^{\lfloor j+1/2 \rfloor} (1 + 4\alpha^2(j+1-n)^2). \quad (7)$$

These results are readily checked for small values of j upon using explicit matrices, say $\hat{\mathbf{n}} \cdot \mathbf{J} = J_3$. Indeed, this is how (6) was first deduced.

Subsequently, after the first version of this paper was posted on the arXiv, a detailed proof of (6) was given in [12]. But, as it turns out, after the previous version of this paper was posted, I learned that (5) and (6) are true for any $N \times N$ matrix M , upon replacing $\hat{\mathbf{n}} \cdot \mathbf{J} \rightarrow M$ and $2j+1 \rightarrow N$. The result is a widely known theorem from the resolvent formalism (e.g. see [13]), and is proven in numerous textbooks and articles on matrix theory (e.g. see [14] and [15]).

Still, some comments are in order. Firstly, note that for either bosonic (integer) or fermionic (semi-integer) spins, only *even* powers of α with positive coefficients are produced by the determinant factors in (6). Consequently the $\mathfrak{A}_k^{[j]}(\alpha)$ and $\mathfrak{B}_k^{[j]}(\alpha)$ coefficients have no singularities for real α . Secondly, the determinants in (6) are essentially generating functions of the central factorial numbers $t(m, n)$ (see [16]), a fact already exploited in [1, 7]. For example, for integer j ,

$$\text{Trunc}_{2j-k}[\det(1 - 2i\alpha \hat{\mathbf{n}} \cdot \mathbf{J})] = \sum_{m=0}^{\lfloor j-k/2 \rfloor} 4^m \alpha^{2m} |t(2j+2, 2j+2-2m)|, \quad (8)$$

with the full determinant obtained for $k = 0$. Thirdly, as $j \rightarrow \infty$ for any fixed k the truncation in (6) is lifted — albeit not without some subtleties [17] — to obtain, for small α , $\lim_{j \rightarrow \infty} \mathfrak{B}_k^{[j]}(\alpha) \sim \alpha^k$. Nevertheless, in contrast to the periodicized θ -monomials found in [1], the large j behavior here does not make the periodicity of rotations manifest. To exhibit periodicity even for finite values of j , θ must be expressed, on a case-by-case basis, as cyclometric functions of α , and then the branch structure of those cyclometric functions must be invoked.

Finally, the Hille-Yosida theorem from resolvent theory provides a one-to-one relation between (2) and (6). That is, for any hermitian $N \times N$ matrix, a Laplace transform gives

$$\frac{1}{1 - itM} = \int_0^\infty e^{-s} e^{istM} ds . \quad (9)$$

So for linearly independent powers M^k , $0 \leq k < N$, as is the case for spin matrices, the matrix polynomial expansion coefficients are related order-by-order.

$$\frac{1}{1 - itM} = \sum_{n=0}^{N-1} A_n(t) (iM)^n , \quad \exp(i\phi M) = \sum_{n=0}^{N-1} C_n(\phi) (iM)^n , \quad A_n(t) = \int_0^\infty e^{-s} C_n(st) ds . \quad (10)$$

This provides yet another route to prove the CFZ result (1) starting from the known coefficients for the resolvent (see [12]). For example, for spin $j = 1/2$,

$$\exp\left(\frac{1}{2} i\phi \sigma_3\right) = \cos\left(\frac{\phi}{2}\right) I + i \sin\left(\frac{\phi}{2}\right) \sigma_3 , \quad (11)$$

$$\int_0^\infty e^{-s} \cos(st/2) ds = \frac{1}{1 + \frac{1}{4} t^2} , \quad \int_0^\infty e^{-s} \sin(st/2) ds = \frac{\frac{1}{2} t}{1 + \frac{1}{4} t^2} , \quad (12)$$

$$\det\left(1 - \frac{1}{2} it \sigma_3\right) = 1 + \frac{1}{4} t^2 , \quad (13)$$

$$\frac{1}{1 - \frac{1}{2} it \sigma_3} = \frac{1}{1 + \frac{1}{4} t^2} \left(I + \frac{1}{2} it \sigma_3\right) , \quad (14)$$

where $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and the Pauli matrix is $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, as usual.

Acknowledgements I thank Curtright and Zachos for suggesting this problem, and for discussions and encouragement. I also thank Jack and Peggy Nichols for their hospitality. Finally, I have benefited from stimulating views of sea waves approaching the Malecón, La Habana.

Keywords: Cayley transform, resolvent, rotations, spin, su(2)

MSC: 15A, 15B, 20C, 22E70, 81R05, 81R50

References

- [1] T L Curtright, D B Fairlie, and C K Zachos, “A compact formula for rotations as spin matrix polynomials” SIGMA 10 (2014) 084. e-Print: arXiv:1402.3541 [math-ph]

- [2] E P Wigner, *Group Theory and its Application to the Quantum Mechanics of Atomic Spectra*, Academic Press (1959).
- [3] Y Lehrer-Ilamed, “On the direct calculations of the representations of the three-dimensional pure rotation group” Proc.Camb.Phil.Soc. 60 (1964) 61–66. DOI: 10.1017/S0305004100037452
- [4] R van Wageningen, “Explicit polynomial expressions for finite rotation operators” Nuc.Phys. 60 (1964) 250–263. DOI: 10.1016/0029-5582(64)90660-1
- [5] T A Weber and S A Williams, “Spin-matrix polynomials and the rotation operator for arbitrary spin” J.Math.Phys. 6 (1965) 1980–1983. DOI: 10.1063/1.1704751
- [6] A J Torruella, “Representations of the three-dimensional rotation group by the direct method” J.Math.Phys. 16 (1975) 1637–1642. DOI: 10.1063/1.522718
- [7] T L Curtright and T S Van Kortryk, “On rotations as spin matrix polynomials” J. Phys. A: Math. Theor. 48 (2015) 025202. e-Print: arXiv:1408.0767 [math-ph]
- [8] *The Collected Mathematical Papers of Arthur Cayley*, Cambridge University Press (1889). See Vol. I. for the Cayley transform, and see Vol. II. for the Cayley-Hamilton theorem.
- [9] See http://en.wikipedia.org/wiki/Cayley_transform. For a recent paper which provides a brief guide to more current literature and some applications, see Y Hardy, A Fošner, and W-H Steeb, “Cayley transform and the Kronecker product of Hermitian matrices” Linear Algebra and its Applications 439 (2013) 4023–4031. DOI: 10.1016/j.laa.2013.09.043
- [10] See http://en.wikipedia.org/wiki/Cayley-Hamilton_theorem.
- [11] In this paper I only use Cayley transforms to construct a group element U from an element S in the Lie algebra, i.e. $U = \frac{1+iS}{1-iS}$. I do not consider the inverse problem to construct the Lie algebra generator from a given group element, i.e. $iS = \frac{U-1}{U+1}$.
- [12] T L Curtright, “More on Rotations as Spin Matrix Polynomials” e-Print: arXiv:1506.04648 [math-ph]
- [13] M X He and P E Ricci, “On Taylor’s formula for the resolvent of a complex matrix” Computers and Mathematics with Applications 56 (2008) 2285–2288. DOI: 10.1016/j.camwa.2008.03.051
- [14] A S Householder, *The Theory of Matrices in Numerical Analysis*, Dover (1975). See Section 6.7, and also the references cited therein, where the history of the identity is traced back to U J J Leverrier, “Sur les variations seculaires des elements des orbites” J. Math. (1840).
- [15] S-H Hou, “A Simple Proof of the Leverrier-Faddeev Characteristic Polynomial Algorithm” SIAM Review 40 (1998) 706–709.
- [16] P L Butzer, M Schmidt, E L Stark, and L Vogt, “Central Factorial Numbers: Their main properties and some applications” Numerical Functional Analysis and Optimization 10 (1989) 419–488.
- [17] The subtleties involved in the limit of large j are discussed more fully in [12] where the precise meaning of $\lim_{j \rightarrow \infty} \mathfrak{B}_k^{[j]}(\alpha) \sim \alpha^k$ is explained in detail, and where exact $\lim_{j \rightarrow \infty} \mathfrak{B}_k^{[j]}(\alpha)$ results are obtained for both integer and semi-integer j .